

Specific Heat of a Three Dimensional Metal Near a $T = 0$ Magnetic Transition with Dynamic Exponent $z = 2, 3$, or 4

U. Zülicke and A. J. Millis

AT&T Bell Laboratories, 600 Mountain Ave, Murray Hill, NJ 07974, USA

Abstract

We derive expressions for the universal contribution to the specific heat of a three-dimensional metal near a zero-temperature phase transition with dynamic exponent $z = 2, 3$, or 4 . The results allow a quantitative comparison of theory to data. We illustrate the application of our results by analyzing data for $\text{Ce}_{1-x}\text{Lu}_x\text{Cu}_2\text{Si}_2$, which has been claimed to be near a quantum critical point.

I. INTRODUCTION

Many strongly-correlated metallic compounds [1] exhibit "non-fermi-liquid" behavior in the sense that the low temperature (T) specific heat (C) exhibits more rapid temperature dependence than expected from the standard expression for a fermi liquid with fermi temperature T_F , which is

$$\frac{C}{T} = \frac{A}{T_F} \left(1 + B \frac{T^2}{T_F^2} \log T + \mathcal{O}\left(\frac{T^2}{T_F^2}\right) + \dots \right) \quad (1.1)$$

with $A = \frac{k_B \pi^2}{2} n$ (n denotes the electronic density) and B given in terms of Landau parameters [2]. Many authors have suggested that the non-fermi-liquid behavior is due to the proximity of the non-fermi-liquid materials to a zero temperature critical point [3–6]. One class of proposed critical points would separate a $T = 0$ magnetically ordered phase from a phase with no long range order [3–5]; another class involves critical points occurring in models of isolated impurities [6]. To experimentally distinguish these intriguing suggestions it is necessary to compare the data to theory. In this communication we provide quantitative expressions for the universal contributions to the specific heat of a three dimensional metal near a zero temperature magnetic—non-magnetic phase transition with a dynamical exponent $z = 2, 3$, or 4 . The case $z = 2$ describes antiferromagnetic transitions not driven by $2k_F$ instabilities or nesting; the case $z = 3$ describes ferromagnetic transitions in a clean metal and also describes the recently discussed "gauge theories" [7] of strongly correlated systems; the case $z = 4$ would describe a dirty ferromagnet in the temperature regime in which the randomness leads to spin diffusion but does not otherwise change the critical properties.

Our results are obtained by solving scaling equations derived previously [3]. The scaling equations imply that the critical behavior is in the Gaussian universality class, with a dangerously irrelevant operator. The results depend on three dimensionless parameters: reduced temperature $t = \frac{T}{T^*}$ (where T^* is a microscopic temperature presumably of the order of the fermi temperature T_F), a control parameter r (which depends on temperature and on a Hamiltonian parameter such as pressure or doping whose variation tunes the material

through the $T = 0$ transition, which is defined to occur at $r(T = 0) = 0$ and u , the coefficient of the "dangerously irrelevant operator". The results of course also depend on the spatial dimensionality d and the dynamical exponent z , and the results we write are only valid for $d + z > 4$. A qualitative phase diagram is shown in Fig.1. The axes are reduced temperature, t , and the zero-temperature control parameter $\delta = r(T = 0)$. A phase transition line $t_c(\delta)$ separates a magnetically ordered phase (shaded region) from a phase with no long range order. We give results for the specific heat coefficient $\gamma = C/T$ in the disordered phase. The behavior in the ordered phase depends on details of the order. Too many special cases arise to discuss here. In the disordered phase we find that the singular part, γ_{sing} , of $\gamma = \frac{C}{T}$ may be written

$$\gamma_{sing} = \frac{k_B}{T^*} t^{\frac{d-z}{z}} g_{d,z} \left(\frac{r}{t^{2/z}}, \frac{ut^{\frac{d+z-2}{z}}}{r} \right) \quad (1.2)$$

where $g_{d,z}$ is a universal scaling function which we calculate. The leading behavior of γ_{sing} is given by $\frac{k_B}{T^*} t^{\frac{d-z}{z}} g_{d,z} \left(\frac{r}{t^{2/z}}, 0 \right)$ and may be computed directly from the Gaussian model. The function $g_{d,z} \left(\frac{r}{t^{2/z}}, 0 \right)$ describes the crossover between a low-temperature (so-called quantum) regime with $t \ll r^{\frac{z}{2}}$ and a high-temperature (so-called classical) regime defined by $t \gg r^{\frac{z}{2}}$. A generally valid expression for $g_{3,z}(x, 0)$ is given in Eq. (B3). Expansions for the quantum and classical limits are given in Eqs. (3.1)–(3.6). We also investigate the dependence of $g_{d,z}(x, y)$ on its second argument and show that it leads to "corrections to scaling", which are then given as a power series in u (times appropriate combinations of t and r). The only important correction to scaling arises from the nonanalyticity in $g_{d,z}(x, y)$ occurring at $y = \infty$, i.e. $r = 0$. The equation $r(t, \delta, u) = 0$ defines the true transition curve $t_c(\delta)$. A precise expression is given in Eq. (3.7). To an extremely good approximation, this transition is a second order mean field transition and is characterized by a specific heat jump which we calculate. The result is Eq. (3.8). We also calculate the Gaussian fluctuation correction to this specific heat jump. This is given in Eq. (3.9).

The rest of this paper is organized as follows. In Section II we write down the model to be solved and indicate the method of solution. In Section III we give the results for $d = 3$

and $z = 2, 3, 4$. In Section IV we illustrate the comparison of the results to data [8] by comparing $g_{3,2}$ to data on $\text{Ce}_{1-x}\text{Lu}_x\text{Cu}_2\text{Si}_2$, a three-dimensional material which is argued to be near a $T = 0$ antiferromagnetic transition. Section V is a brief conclusion. Details of various calculations are given in Appendices.

II. FORMALISM

We follow the previous approach [3,10] of integrating out the fermionic degrees of freedom and writing the partition function as a functional integral over an N-component bosonic order parameter $\vec{\phi}$ which represents the ordering field. This procedure is not useful for superconducting transitions, transitions at nesting wavevectors of nested fermi surfaces, or indeed transitions at wavevectors Q connecting two points on the fermi surface with parallel tangents (e.g. $Q = 2k_F$ for a spherical fermi surface) because the resulting bosonic action is singular. For further details see Refs. [3,9,10]. Recently, Sachdev [11] has questioned whether the action used in Refs. [3,10] is adequate to describe magnetic phase transitions in which the order parameter has $O(3)$ or higher symmetry because of the neglect of terms giving rise to the precession terms in the spin-wave equation of motion, but it has been recently argued that the action of Refs. [3,10] is in fact correct even in this case [12]. We do not discuss this issue here, but assume the validity of the approach of Refs. [3,10], which is certainly correct for low symmetry order parameters and may apply to higher symmetries.

The partition function is

$$\mathcal{Z} = \int \mathcal{D}\vec{\phi} \exp(-\mathcal{S}[\vec{\phi}]) \quad (2.1)$$

and the action $\mathcal{S}[\vec{\phi}]$ is given by

$$\mathcal{S}[\vec{\phi}] = \mathcal{S}^{(0)} + \mathcal{S}^{(2)}[\vec{\phi}] + \mathcal{S}^{(4)}[\vec{\phi}] + \dots \quad (2.2)$$

$$\mathcal{S}^{(2)}[\vec{\phi}] = V \sum_n \int_{\xi_0}^{\Lambda} \frac{d^d k}{(2\pi)^d} \left[\delta_0 + (\xi_0 k)^2 + \frac{|\Omega_n|}{\Upsilon(\xi_0 k)^{z-2}} \right] \vec{\phi}_n(\vec{k}) \vec{\phi}_{-n}(-\vec{k}) \quad (2.3)$$

$$\mathcal{S}^{(4)}[\vec{\phi}] = u_0 V^3 \sum_{n_1, \dots, n_4} \int_{\xi_0}^{\Lambda} \frac{d^d k_1}{(2\pi)^d} \dots \frac{d^d k_4}{(2\pi)^d} \vec{\phi}_{n_1}(\vec{k}_1) \dots \vec{\phi}_{n_4}(\vec{k}_4) \delta(\sum_i \vec{k}_i) \delta_{\sum_i n_i} \quad (2.4)$$

Here $\mathcal{S}^{(0)}$ is the logarithm of the contribution to the partition function from the degrees of freedom which have been integrated out. It depends on T and other external parameters. The parameter δ_0 tunes the system through the zero-temperature critical point and might be related to the pressure or to chemical composition. The coupling strength u_0 is assumed to be constant near the critical point. The relation of δ_0 to the above mentioned control parameter r will be given below. The wavevector k is measured from the ordering wave vector Q . ξ_0 is a microscopic length, presumably of the order of k_F^{-1} . A microscopic energy scale is given by Υ which one expects to be of the order of E_F . The summation index n labels the Matsubara frequencies $\Omega_n = 2\pi n k_B T$. Momentum integrals are cut off at $\frac{\Lambda}{\xi_0}$, energy integrals obtained by analytically continuing Matsubara sums are cut off at $\Gamma\Upsilon(\xi_0 k)^{z-2}$. In a fermi liquid one expects the dimensionless cutoff parameters Λ and Γ to be of order unity. The dots indicate terms which are irrelevant at the critical point.

In Ref. [3] the critical behavior of the model was shown to be determined by perturbative (in u_0) scaling about the $u_0 = 0$ (Gaussian) fixed point. The scaling procedure used in [3] maps $\mathcal{Z}$ onto a new $\mathcal{Z}'$ identical in form to $\mathcal{Z}$ but with rescaled parameters $T(b), \delta(b), u(b)$. The solution of the scaling equations can be written

$$t(b) = tb^z \quad (2.5)$$

$$u(b) = ub^{4-(d+z)} + \mathcal{O}(u^2) \quad (2.6)$$

$$\delta(b) = b^2 \left\{ \delta_0 + 2u(N+2) \int_0^{\log b} dx e^{[2-(d+z)]x} f^{(2)}(te^{zx}, \delta(e^x)) \right\} + \mathcal{O}(u^2) \quad (2.7)$$

$$t \mathcal{S}^{(0)}(b) = \frac{V}{\xi_0^d} N \int_0^{\log b} dx e^{-(d+z)x} f^{(0)}(te^{zx}, \delta(e^x)) + S_0^{(0)} \quad (2.8)$$

Here $t = \frac{T}{T^*}$ with $T^* = \frac{\Upsilon}{k_B}$ and $u = \frac{u_0}{u^*}$ with $u^* = \frac{V^3}{\xi_0^{3d}}$. The analytical expressions for $f^{(0)}(x, y)$ and $f^{(2)}(x, y)$ have been obtained in [3] and read:

$$\begin{aligned} f^{(0)}(x, y) &= -\frac{K_d \Lambda^{d+z-2}}{(2\pi)^d} \int_0^\Gamma \frac{d\varepsilon}{\pi} \coth\left(\frac{\Lambda^{z-2}\varepsilon}{2x}\right) \arctan\left(\frac{\varepsilon}{y + \Lambda^2}\right) \\ &\quad - \frac{2}{\pi} \Gamma \int_0^\Lambda \frac{d^d k}{(2\pi)^d} k^{z-2} \coth\left(\Gamma \frac{k^{z-2}}{2x}\right) \arctan\left(\frac{\Gamma}{y + k^2}\right) \\ f^{(2)}(x, y) &= \frac{K_d \Lambda^{d+z-2}}{(2\pi)^d} \int_0^\Gamma \frac{d\varepsilon}{\pi} \coth\left(\frac{\Lambda^{z-2}\varepsilon}{2x}\right) \frac{\varepsilon}{\varepsilon^2 + (y + \Lambda^2)^2} \end{aligned} \quad (2.9)$$

$$+\frac{2}{\pi}\Gamma^2\int_0^\Lambda\frac{d^dk}{(2\pi)^d}\coth(\Gamma\frac{k^{z-2}}{2x})\frac{k^{z-2}}{\Gamma^2+(y+k^2)^2}\quad (2.10)$$

Note we have included the y -dependence of $f^{(2)}$; this was neglected in Ref. [3]. Its inclusion does not affect physical results in an important way, but makes the mathematical structure more clear.

We turn now to the solution of these equations. For this purpose, we rewrite (2.7)

$$\delta(b) = b^2(r(t, \delta, u) - u \Delta(t, r(t, \delta, u), b)) + \mathcal{O}(u^2) \quad (2.11)$$

with

$$r(t, \delta, u) = \delta + 2u(N+2)\int_0^\infty dx e^{[2-(d+z)]x} [f^{(2)}(te^{zx}, r(t, \delta, u)e^{2x}) - f^{(2)}(0, 0)] \quad (2.12)$$

$$\delta = \delta_0 + 2(N+2)uf^{(2)}(0, 0)/[d+z-2] \quad (2.13)$$

and

$$\Delta(t, r(t, \delta, u), b) = 2(N+2)\int_{\log b}^\infty dx e^{[2-(d+z)]x} f^{(2)}(te^{zx}, r(t, \delta, u)e^{2x}) \quad (2.14)$$

We can neglect the $\mathcal{O}(u^2)$ -terms in (2.11), because they do not affect the leading nonanalytic behavior in u . The expression (2.12) is calculated in Appendix A. The important result for our subsequent considerations in this Section is that we can write $r(t, \delta, u)$ as

$$r(t, \delta, u) = \delta + (N+2)ut^{\frac{d+z-2}{z}}\phi_{d,z}\left(\frac{r}{t^{2/z}}\right) \quad (2.15)$$

with $\phi_{d,z}$ given by Eq. (A6).

To calculate the free energy we use the scaling equation [13]

$$F(t, \delta, u) = b^{-(d+z)}F(tb^z, \delta(b), u(b)) + t\mathcal{S}^{(0)}(b) \quad (2.16)$$

with $F(t, \delta, u)$ being the free energy of a system described by the model (2.2), calculated perturbatively in the parameter u and measured in units of $k_B T^*$. Investigation of the scale (b-) dependent terms up to $\mathcal{O}(u)$ in (2.16) shows that we can write (2.16) as a trajectory integral in phase space [14]:

$$F(t, \delta, u) = \frac{V}{\xi_0^d} N \int_0^\infty dx e^{-(d+z)x} f^{(0)}(te^{zx}, \delta(e^x)) + \mathcal{O}(u^2) \quad (2.17)$$

where the $\mathcal{O}(u^2)$ -terms do not contain the leading nonanalytic behavior in u . Inserting expression (2.11) for $\delta(e^x)$ and performing a variable change we get

$$F(t, \delta, u) = \frac{V}{\xi_0^d} N t^{\frac{d+z}{z}} \int_{\frac{1}{z} \log t}^\infty dx e^{-(d+z)x} f^{(0)} \left(e^{zx}, \frac{r}{t^{\frac{z}{2}}} \left[1 - \frac{ut^{\frac{d+z-2}{z}}}{r} \Delta \left(1, \frac{r}{t^{\frac{z}{2}}}, e^x \right) \right] e^{2x} \right) \quad (2.18)$$

It is therefore apparent that we can write the free energy (apart from non-singular contributions) as

$$F(t, \delta, u) = t^{\frac{d+z}{z}} f \left(\frac{r}{t^{\frac{z}{2}}}, \frac{ut^{\frac{d+z-2}{z}}}{r} \right) \quad (2.19)$$

Due to the relation (2.15) and the definition of $\Delta(t, r(t, \delta, u), b)$, this scaling behavior is preserved when taking derivatives with respect to temperature, and we find the specific heat coefficient to obey (1.2). It is shown in Appendix C that the dependence on the variable $\frac{ut^{\frac{d+z-2}{z}}}{r}$ leads to negligible corrections to the leading scaling behavior. The latter is given by

$$\gamma_{\text{leading}} = \frac{V}{\xi_0^d} N \int_0^\infty dx e^{(z-d)x} \partial_\zeta^2 f^{(0)}(\zeta, r e^{2x}) \Big|_{\zeta=te^{zx}} \quad (2.20)$$

and is evaluated in detail in Appendix B.

In order to characterize the specific heat near the transition line $t_c(\delta)$ we calculate the discontinuity (jump) of the specific heat as it is predicted by mean field theory, using the scaling equation (2.16) for the free energy. Very close to the true transition, the model (2.2) can be scaled to a classical Gaussian model given by the Landau-Ginzburg functional [3]

$$\exp \left(-\frac{F(t(b), \delta(b), u(b))}{t(b)} \right) = \int \mathcal{D}\phi \exp(-\mathcal{S}[\vec{\phi}]) \quad (2.21)$$

$$\mathcal{S}[\vec{\phi}] = \int d^d x \left\{ \frac{1}{2} |\vec{\nabla} \vec{\phi}|^2 + \delta(b) |\vec{\phi}|^2 + v(b) (|\vec{\phi}|^2)^2 \right\} \quad (2.22)$$

with $v(b) = u(b)t(b)$. Applying the usual mean field approximation yields the discontinuity of the specific heat. Analytical details of this calculation are given in Appendix D. The result is Eq. 3.8. The correction to the specific heat jump due to Gaussian fluctuations is

also evaluated in this Appendix. The result is Eq. (3.9). Non-gaussian critical fluctuations become important when the Ginzburg criterion is violated [13]. The Ginzburg criterion may be obtained by demanding that the Gaussian fluctuation contribution to the specific heat is small compared to the mean field specific heat jump. The result is $ut_cr^{\frac{d-4}{2}} \ll 1$. This is violated for $t - t_c < t_c^{2-\frac{1}{z}}$ in 3 dimensions. The expressions for the Ginzburg criterion and the width about the transition line where it is violated correct the results obtained in [3], which were in error by factors of $t_c^{1/z}$. As the critical temperature is small, mean field behavior characterizes the transition correctly except from a negligibly small region about t_c .

III. RESULTS FOR 3D

We give the results for the specific heat coefficient (normalized per mol) and for the transition temperature in the physically relevant case of 3 dimensions in this section.

The formula describing the specific heat in the whole Gaussian region of the phase diagram is Eq. (B3) [which can be simplified to (B4) for $z = 2$]. However, in order to compare experimental data to the prediction of this formula it is necessary to extract the system-dependent scale T^* and the scale factor which relates $r(T = 0)$ to experimentally accessible control parameters like pressure or concentration first. For this purpose it is useful to have expressions for γ in the quantum ($T \ll T^*r$) and classical ($T \gg T^*r$) limits in the phase diagram. Fits of the data to the appropriate functional forms provide the scales which can then be inserted into (B3), which must in general be evaluated numerically.

The scale factors r^* (which we have set equal to 1) and T^* are of course arbitrary. Only dimensionless combinations of the coefficients we present are universal. However, there is a natural definition of the scale factor which gives an estimate of the important microscopic scales of the system. We have defined T^* so that for $d = 3$ and $z = 2$ the leading low- T non-singular contribution to the Gaussian model specific heat is $\frac{k_B}{T^*} \frac{\Lambda}{\xi_0^d} \frac{N}{3\pi}$. For a density of order 1, $\xi_0^{-1} \sim k_F \sim (3\pi^2)^{\frac{1}{3}}$ so the specific heat coefficient would be of the order of the

leading fermi-liquid term in (1.1).

Furthermore, it is important to note that the specific heat calculated in this communication is only the contribution arising from the existence of a quantum critical point. A metallic system has an additional background specific heat given by (1.1). Depending on the dimensionality and the dynamical exponent, the leading behavior or corrections to the leading behavior in (1.1) are changed due to critical fluctuations.

Antiferromagnetism ($z = 2$): The leading term of γ is still the normal fermi liquid contribution, a nonuniversal constant γ_0 . Only corrections to this leading behavior are affected by the critical point. In the quantum regime ($T \ll T^*r$) we find:

$$\gamma^{quantum,z=2} = \gamma_0 - \frac{k_B N_A}{T^*} \frac{N}{6} r^{\frac{1}{2}} - \frac{k_B N_A}{(T^*)^3} \frac{N\pi^2}{60} \frac{T^2}{r^{\frac{3}{2}}} + \dots \quad (3.1)$$

Thus the coefficient of the first ($\mathcal{O}(T^2)$) correction to the fermi liquid behavior diverges as $r \rightarrow 0$ in the quantum regime. The classical limit ($T \gg T^*r$) yields

$$\gamma^{classical,z=2} = \gamma_0 - \frac{k_B N_A}{(T^*)^{\frac{3}{2}}} \frac{15N}{64} \left(\frac{2}{\pi}\right)^{\frac{3}{2}} \zeta\left(\frac{5}{2}\right) T^{\frac{1}{2}} - \frac{k_B N_A}{(T^*)^{\frac{1}{2}}} \frac{3N}{124} \left(\frac{2}{\pi}\right)^{\frac{3}{2}} \zeta\left(\frac{3}{2}\right) \frac{r}{T^{\frac{1}{2}}} + \dots \quad (3.2)$$

with $\zeta(\frac{5}{2}) = 1.34149$ and $\zeta(\frac{3}{2}) = 2.61238$. The term linear in r dominates corresponding terms in the non-critical part of the specific heat because it is enhanced by a factor $T^{-\frac{1}{2}}$ at low temperatures.

Clean Ferromagnet ($z = 3$): In the quantum regime both the leading (constant) term and the $T^2 \log T$ -correction of the normal fermi liquid expression (1.1) are enhanced by critical fluctuations:

$$\gamma^{quantum,z=3} = \gamma_0 - \frac{k_B N_A}{T^*} \frac{N}{6\pi} \log r - \frac{k_B N_A}{(T^*)^3} \frac{2\pi N}{15} \frac{T^2}{r^3} \log \left(\frac{(T/T^*)^2}{r^3} \right) + \dots \quad (3.3)$$

The classical regime shows non-fermi-liquid behavior:

$$\gamma^{classical,z=3} = -\frac{k_B N_A}{T^*} \frac{N}{9\pi} \log \left(\frac{T}{\tilde{T}} \right) - \frac{k_B N_A}{(T^*)^{\frac{1}{3}}} \frac{N\Gamma(\frac{7}{3})\zeta(\frac{4}{3})}{12\pi^2\sqrt{3}} \frac{r}{T^{\frac{2}{3}}} + \dots \quad (3.4)$$

with $\Gamma(\frac{7}{3})\zeta(\frac{4}{3}) = 4.28742$.

Dirty Ferromagnet ($z = 4$): In this case, the leading behavior in the quantum regime is of fermi liquid type and enhanced by critical fluctuations, but the first correction is not

fermi liquid like, and the coefficient of the latter diverges when approaching the critical point.

$$\gamma^{quantum,z=4} = \frac{k_B N_A}{T^*} \frac{N}{6} r^{-\frac{1}{2}} - \frac{k_B N_A}{(T^*)^{\frac{3}{2}}} \frac{15N}{64} \left(\frac{2}{\pi}\right)^{\frac{3}{2}} \zeta\left(\frac{5}{2}\right) \frac{T^{\frac{1}{2}}}{r^{\frac{3}{2}}} + \dots \quad (3.5)$$

In the classical regime, the leading behavior is determined by critical fluctuations and is not of fermi liquid type:

$$\gamma^{classical,z=4} = -\frac{k_B N_A}{(T^*)^{\frac{3}{4}}} \frac{N}{8\pi^2} \frac{\Gamma(\frac{11}{4})\zeta(\frac{7}{4})}{\sin(\frac{7}{8}\pi)} T^{-\frac{1}{4}} - \frac{k_B N_A}{(T^*)^{\frac{1}{4}}} \frac{N}{32\pi^2} \frac{\Gamma(\frac{9}{4})\zeta(\frac{5}{4})}{\sin(\frac{3}{8}\pi)} \frac{r}{T^{\frac{3}{4}}} + \dots \quad (3.6)$$

with $\Gamma(\frac{11}{4})\zeta(\frac{7}{4}) = 3.15612$ and $\Gamma(\frac{9}{4})\zeta(\frac{5}{4}) = 5.20628$.

Transition region: In order to characterize the region near the transition line $T_c(\delta)$ we give the result for this curve in 3D

$$T_c = T^* \left[\frac{-\delta}{\mathcal{D}_{3,z}(N+2)u} \right]^{\frac{z}{1+z}} \quad (3.7)$$

the expression for the specific heat jump predicted by mean field theory

$$\Delta C = \frac{k_B N_A}{(T^*)^{\frac{2}{z}(z+1)}} N(N+2)^2 \mathcal{D}_{3,z}^2 u T_c^{\frac{2+z}{z}} \quad (3.8)$$

and the correction to the mean field result due to Gaussian fluctuations

$$C^{transition} = k_B N_A \frac{N}{16\pi} \left(\frac{z}{z+1}\right)^{\frac{1}{2}} \left[\mathcal{D}_{3,z}(N+2)u \left(\frac{T_c}{T^*}\right)^{\frac{z+1}{z}} \right]^{\frac{3}{2}} \left[\frac{T - T_c}{T_c} \right]^{-\frac{1}{2}} \quad (3.9)$$

The numerical values for $\mathcal{D}_{3,z}$ are $\mathcal{D}_{3,2} = 0.52109$, $\mathcal{D}_{3,3} = 0.39396$, and $\mathcal{D}_{3,4} = 0.35875$.

IV. COMPARISON TO EXPERIMENTAL DATA

The discovery of superconductivity in so-called heavy-fermion systems [1] resulted in extensive studies of these materials. Some of them undergo magnetic phase transitions at low temperatures, others display no long-range order. It has become evident, however, that several of the latter are near magnetic critical points as well. As an example, modest doping of CeCu₆ [16,17] and UPt₃ [18,19] leads to antiferromagnetism. The influence of pressure variation on the critical behavior has been studied as well [17,20]. The reason for the

proximity of the heavy-fermion metals to a quantum critical point is the extreme sensitivity of the two competing energy scales (fermi [Kondo] temperature for the fermi liquid state and RKKY exchange energy for the magnetically ordered state) in these materials to changes in parameters such as the lattice constant. Therefore the heavy fermion compounds may be an appropriate experimental system to study the implications of the theory presented in this communication.

Application to CeCu₂Si₂

Measurements of the temperature-dependence of the specific heat have been carried out at compounds Ce_{1-x}M_xCu₂Si₂ with M=La,Y,Lu and $x = 0.0, 0.05, 0.1, 0.2, 0.5$ and 0.9 [8]. Doping with La expands the lattice, whereas doping with either Y or Lu contracts it. As the data for La-doped samples show a rather complicated behavior, we will not comment on them in this work. Diluting CeCu₂Si₂ with Y,Lu indicates, however, that this material is near a quantum-critical point. Based on the assumption that this is an antiferromagnetic instability, we present an interpretation of the data for Ce_{1-x}Lu_xCu₂Si₂. The numbers and conclusions which can be drawn from the experiments carried out with Y-doped samples do not differ significantly from those given below.

In Fig.2 the data for various concentrations x are shown. The parameter tuning the system through the quantum critical point $\delta = r(T = 0)$ is defined as $\delta = \frac{x-x_c}{x^*}$. If the pure compound is right at the critical point ($x_c = 0$), this system would be in the classical regime and we would expect to find the low-temperature specific heat behaving according to (3.2). Indeed the low-temperature ($T \leq 5K$) data for $x = 0$ fit nicely to the square root temperature-dependence. This fit yields the temperature scale $T^* = 5K$ which is of the order of the Kondo temperature in this material: $T_K = 10K$ [21]. Furthermore, the first correction to the leading behavior in the classical regime is linear in r and therefore linear in δ as well. As we will check below, the compounds for $x = 0.05, 0.1$ are still in the classical regime. We take the measured values for $\frac{C}{T}(T = 1.1K)$ given in [8] at concentrations $x = 0.0$ and $x = 0.05$ in order to determine the scale x^* . We find $x^* = 0.3$, hence for the

quantum-classical crossover temperature $T_{cross} = \frac{T^*}{x^*} x = 17x$ which is reached right at the beginning of the measurements for $x = 0.05, 0.1$, whereas for $x = 0.2$ the quantum-classical crossover is accessed experimentally. The data for $x = 0.5, 0.9$ and $T > 5K$ cannot be explained within the framework of the theory presented here because the systems are already too far away from the critical point. Taking the values for the nonuniversal scale factors $T^* = 5K$, $x^* = 0.3$, we can apply (B4) to predict the temperature dependence of the specific heat for $x = 0.05, 0.1, 0.2$. As Fig.2 shows, theory and data agree quite well for low temperatures. Please note that we neglected any effect of the dangerously irrelevant variable u . The coincidence for $x = 0.2$ at low temperatures is particularly nice because there the system is in the quantum regime, whereas we extracted all information for the theory from the classical regime. There is apparently a systematic deviation from the theory occurring at lower temperatures for larger x . We do not understand this deviation at present. We suspect that the deviations (starting at lower temperatures for increasing concentration) are due to noncritical fluctuations which become important farther away from the critical point, but the observed interplay between T and x is not expected in our model.

V. CONCLUSION

We have determined the universal functions needed to quantitatively analyze the behavior of the specific heat near a quantum critical point with dynamical exponent $z = 2, 3, 4$ in $d = 3$ spatial dimensions and illustrated, by an example, the use of the equations. The results are summarized in Eqs. (3.1)–(3.9), (B3) and (B4), respectively. Depending on the dynamical exponent z , the leading or subleading behavior of the normal fermi liquid specific heat coefficient is affected due to the proximity to the quantum critical point.

The physical properties of these transitions are governed by four non-universal scales: a temperature scale T^* , a scale r^* for the Hamiltonian parameter tuning the $T = 0$ transition, a dangerously irrelevant variable u , and a length scale ξ_0 . Once these are known, all the behavior is fixed. The specific heat does not involve the length scale ξ_0 but provides sufficient

information to overdetermine the remaining parameters.

An important open theoretical problem is to understand the nature and magnitude of the corrections to the universal behavior. On the experimental side, it would be useful to study in detail the specific heat of a system to which the theory applies.

ACKNOWLEDGMENTS

We thank P. C. Hohenberg for helpful discussions. U. Z. was supported in part by the Studienstiftung des deutschen Volkes (German National Scholarship Foundation).

APPENDIX A: QUANTUM PARAMETER AND CRITICAL TEMPERATURE

In this appendix we will calculate (2.12) and obtain an expression for the critical temperature.

We first extract the explicit temperature dependence of r when writing

$$r(t, \delta, u) = \tilde{r}(r(t, \delta, u), u) + \tilde{\tilde{r}}(t, r(t, \delta, u), u) \quad (\text{A1})$$

$$\tilde{r}(\delta, u) = \delta + 2u(N+2) \int_0^\infty dx e^{[2-(d+z)]x} \left[f^{(2)}(0, r(t, \delta, u) e^{2x}) - f^{(2)}(0, 0) \right] \quad (\text{A2})$$

$$\tilde{\tilde{r}}(t, \delta, u) = 2u(N+2) \int_0^\infty dx e^{[2-(d+z)]x} \left[f^{(2)}(te^{zx}, r(t, \delta, u) e^{2x}) - f^{(2)}(0, r(t, \delta, u) e^{2x}) \right] \quad (\text{A3})$$

Further inspection of Eq. (A2) reveals that

$$\tilde{r}(\delta, u) = \delta \left(1 + 2u(N+2) \frac{\partial_\eta f^{(2)}(0, \eta)|_{\eta=0}}{d+z-4} \right) + 2u(N+2) \begin{cases} \mathcal{O}(r^{\frac{d+z-2}{2}}) & \text{if } d+z-6 \neq 0 \\ \mathcal{O}(r^2 \log r) & \text{if } d+z-6 = 0 \end{cases} \quad (\text{A4})$$

where we omitted $\mathcal{O}(u^2)$ -terms like those we have already neglected when writing (2.11). The dangerously irrelevant variable u leads to a rescaling of δ by a u -dependent factor, which can be absorbed into the scale (unit of measurement) of δ . Note that the nonanalytic contributions in r due to the variable u are nonleading and therefore negligible near the critical point.

Neglecting nonleading (analytic) contributions we can write for (A3)

$$\tilde{\tilde{r}}(t, \delta, u) = u(N+2) t^{\frac{d+z-2}{z}} \phi_{d,z} \left(\frac{r}{t^{2/z}} \right) + \mathcal{O}(ut^2) \quad (\text{A5})$$

with

$$\phi_{d,z}(x) = \frac{K_d}{(2\pi)^{d+1}} 8 \int_0^\infty dy y^{\frac{z-d}{2}} \int_0^\infty d\varepsilon \frac{\varepsilon(\coth \varepsilon - 1)}{4\varepsilon^2 y^z + (xy + 1)^2} \quad (\text{A6})$$

The two limits $\frac{r}{t^{2/z}} \ll 1$ and $\frac{r}{t^{2/z}} \gg 1$ correspond to the classical and quantum regimes, respectively [3]. We find

$$\begin{aligned} \phi_{d,z}(x) &\Rightarrow \mathcal{D}_{d,z} && \text{for } x \rightarrow 0 \text{ (classical)} \\ \phi_{d,z}(x) &\Rightarrow x^{\frac{d-z-2}{2}} && \text{for } x \rightarrow \infty \text{ (quantum)} \end{aligned}$$

with

$$\mathcal{D}_{d,z} = \frac{K_d}{(2\pi)^d} \frac{2}{z} \frac{\Gamma(\frac{d-2}{z} + 1) \zeta(\frac{d-2}{z} + 1)}{\sin \frac{z+2-d}{2z} \pi} \quad (\text{A7})$$

Numerical values for $\mathcal{D}_{d,z}$ in three dimensions and for the dynamical exponents considered in this communication are $\mathcal{D}_{3,2} = 0.52109$, $\mathcal{D}_{3,3} = 0.39396$, and $\mathcal{D}_{3,4} = 0.35875$. The term with explicit temperature dependence in the quantum regime is $t^2 r^{\frac{d-z-2}{2}}$ which is small compared to the contribution $r^{\frac{d+z-2}{2}}$ to $\tilde{r}$ which we have already neglected. The general result is therefore

$$r(t, \delta, u) = \delta + (N+2) u t^{\frac{d+z-2}{z}} \phi_{d,z} \left(\frac{r}{t^{2/z}} \right) \quad (\text{A8})$$

$$= \delta \text{ in the quantum regime} \quad (\text{A9})$$

$$= \delta + \mathcal{D}_{d,z} (N+2) u t^{\frac{d+z-2}{z}} \text{ in the classical regime} \quad (\text{A10})$$

The true transition occurs when r vanishes. Therefore we find an expression for the transition temperature t_c by setting (A10) equal to zero and solving for t :

$$t_c = \left[\frac{-\delta}{\mathcal{D}_{d,z} (N+2) u} \right]^{\frac{z}{d+z-2}} \quad (\text{A11})$$

APPENDIX B: QUANTUM-CLASSICAL CROSSOVER

In this section we investigate the expression (2.20) for the leading contributions to the specific heat coefficient, which arise from the quantum-classical crossover.

The expression (2.20) can be written

$$\gamma_{\text{leading}} = \frac{V}{\xi_0^d} \frac{4K_d N}{(2\pi)^{d+1}} \int_{\frac{1}{\Lambda^2}}^{\infty} dx x^{\frac{z-d}{2}-1} \int_0^{\infty} d\varepsilon \frac{\varepsilon^2}{\sinh^2 \varepsilon} \frac{rx + 1}{(2t\varepsilon)^2 x^z + (rx + 1)^2} \quad (\text{B1})$$

$$= r^{\frac{d-z}{2}} \frac{V}{\xi_0^d} \frac{4K_d N}{(2\pi)^{d+1}} \int_0^{\infty} d\varepsilon \frac{\varepsilon^2}{\sinh^2 \varepsilon} \int_{\frac{r}{\Lambda^2}}^{\infty} dx x^{\frac{z-d}{2}-1} \frac{x + 1}{(x + 1)^2 + \frac{(2t\varepsilon)^2}{r^z} x^z} \quad (\text{B2})$$

$$= t^{\frac{d-z}{z}} \frac{V}{\xi_0^d} \frac{2^{\frac{d+z}{z}} K_d N}{(2\pi)^{d+1}} \int_0^{\infty} d\varepsilon \frac{\varepsilon^{\frac{d+z}{z}}}{\sinh^2 \varepsilon} \int_{\frac{(2t\varepsilon)^{2/z}}{\Lambda^2}}^{\infty} dx x^{\frac{z-d}{2}-1} \frac{\frac{r}{(2t\varepsilon)^{2/z}} x + 1}{\left(\frac{r}{(2t\varepsilon)^{2/z}} x + 1 \right)^2 + x^z} \quad (\text{B3})$$

In the case of $d \neq z$, the dependence on the cut-off Λ gives rise to analytic terms $\mathcal{O}(r, t^2)$ only, which we will neglect in the following. To discuss the case $d = z$, however, the existence

of a lower boundary of the x -integral is essential for technical reasons because logarithms occur. But even in this case, the nonanalytic behavior of the heat capacity does not depend on the cut-off.

For $d = 3, z = 2$ it is possible to perform the x -integrals and we find for the singular part of γ which arises from the quantum-classical crossover:

$$\gamma_{leading}^{d=3,z=2} = -t^{\frac{1}{2}} \frac{V}{\xi_0^3} \frac{N}{\sqrt{2\pi^2}} \int_0^\infty d\varepsilon \frac{\varepsilon^2}{\sinh^2 \varepsilon} \sqrt{\frac{r}{t} + \sqrt{\left(\frac{r}{t}\right)^2 + (2\varepsilon)^2}} \quad (\text{B4})$$

In the next paragraph we study the asymptotic behavior of $\gamma_{leading}$ in the classical and quantum regimes corresponding to the limits $\frac{r}{t^{2/z}} \ll 1$ and $\frac{r}{t^{2/z}} \gg 1$, respectively. For sake of simplicity, we do not write the factor $\frac{V}{\xi_0^d} \frac{K_d N}{(2\pi)^{d+1}}$ explicitly. Results of the following paragraphs have to be multiplied by this term to obtain the correct results.

Classical regime: To investigate the classical limit it is most convenient to use Eq. (B3) and expand in the parameter $\frac{r}{t^{2/z}}$ (which is small in the classical regime). The leading behavior is

$$\gamma_{leading}^{classical, d \neq z} = \frac{2\pi}{z} \frac{\Gamma(\frac{d}{z} + 2) \zeta(\frac{d}{z} + 1)}{\sin\left(\frac{d+z}{2z}\pi\right)} t^{\frac{d-z}{z}} + \dots \quad (\text{B5})$$

in the case $d \neq z$, whereas for $d = z$ we have

$$\gamma_{leading}^{classical, d=z} = -\frac{4\pi^2}{3d} \log t + \dots \quad (\text{B6})$$

The first correction to the leading term is obtained by an expansion in the small parameter and reads

$$-\frac{\pi}{2z} \frac{\Gamma(\frac{d-2}{z} + 2) \zeta(\frac{d-2}{z} + 1)}{\sin\left(\frac{2+z-d}{2z}\pi\right)} \frac{r}{t^{\frac{2+z-d}{z}}} \quad (\text{B7})$$

Quantum regime: In this limit we use an expansion of (B2) in the small parameter $\frac{t^2}{r^z}$. The leading term for $d \neq z$ is

$$\gamma_{leading}^{quantum, d \neq z} = \frac{2}{3} \pi^3 \operatorname{sgn}(z - d) r^{\frac{d-z}{2}} + \dots \quad (\text{B8})$$

In the case $d = z$ we find

$$\gamma_{leading}^{quantum,d=z} = -\frac{2}{3}\pi^2 \log \frac{r}{\Lambda^2} + \dots \quad (\text{B9})$$

For $z = 2$ the corrections to the leading behavior are obtained by expanding (B2) in $\frac{t^2}{r^z}$.

This yields the asymptotic series

$$\gamma_{leading}^{quantum,d \neq z=2} = -4\pi^3 r^{\frac{1}{2}} \sum_{n=0}^{\infty} (-1)^n (2\pi)^n |B_{2n+2}| \frac{\Gamma(2n - \frac{1}{2})}{\Gamma(-\frac{1}{2})\Gamma(2n+1)} \left(\frac{t}{r}\right)^{2n} \quad (\text{B10})$$

$$= -\frac{2\pi^3}{3} r^{\frac{1}{2}} - \frac{\pi^5}{15} \frac{t^2}{r^{\frac{3}{2}}} + \dots \quad (\text{B11})$$

where B_m denote Bernoulli numbers. Such an expansion is inappropriate to find corrections in the cases $z = 3, 4$ because these are nonanalytic in the quantity $\frac{t^2}{r^z}$. It is more convenient to rewrite (B3) for large $\frac{r}{t^{2/z}}$ and subtract the already calculated leading term. The corrections are then given by

$$- \frac{2^{\frac{d}{z-2}+1}}{z-2} \frac{t^{\frac{d}{z-2}-1}}{r^{\frac{d}{z-2}}} \int_0^{\infty} d\varepsilon \frac{\varepsilon^{\frac{d}{z-2}+1}}{\sinh^2 \varepsilon} \int_{\frac{(2t\varepsilon)^{2/z}}{r^z}}^{\infty} dx \frac{x^{\frac{z-d-2}{2(z-2)}}}{1+x} \quad (\text{B12})$$

which gives for $d = z = 3$

$$\frac{8}{15}\pi^4 \frac{t^2}{r^3} \log \frac{t^2}{r^3} + \dots \quad (\text{B13})$$

and $d = 3, z = 4$

$$- \frac{15}{2} \left(\frac{\pi}{2}\right)^{\frac{3}{2}} \zeta\left(\frac{5}{2}\right) \frac{t^{\frac{1}{2}}}{r^{\frac{3}{2}}} + \dots \quad (\text{B14})$$

APPENDIX C: DANGEROUSLY IRRELEVANT VARIABLE AND THE QUANTUM-CLASSICAL CROSSOVER

In this appendix we investigate the dependence of (2.18) on the argument $\frac{ut^{\frac{d+z-2}{z}}}{r}$ in the quantum and classical regimes. We show that it leads to corrections to the leading behavior given by (1.2), which is calculated in Appendix B.

In the quantum regime ($t \ll r^{z/2}$) we can write (2.18) as

$$F(t, \delta, u) = \frac{V}{\xi_0^d} N r^{\frac{d+z}{2}} \int_{\frac{1}{2} \log r}^{\infty} dx e^{-(d+z)x} f^{(0)} \left(\frac{t}{r^{z/2}} e^{zx}, \left[1 - ur^{\frac{d+z-4}{2}} \Delta \left(\frac{t}{r^{z/2}}, 1, e^x \right) \right] e^{2x} \right) \quad (\text{C1})$$

As the parameter $\frac{t}{r^{z/2}}$ is small, an expansion in this quantity is reasonable. The term $ur^{\frac{d+z-4}{2}}$ is small compared to unity near the critical point (u is assumed to be of order unity, and r is small), it leads to corrections to coefficients in the leading behavior in the quantum regime, the latter is given by an expansion of (B2) in the small parameter $\frac{t}{r^{z/2}}$.

Similarly, in the classical regime we can expand (2.18) in the small quantity $\frac{r}{t^{2/z}}$. The correction due to the argument $\frac{ut^{\frac{d+z-2}{z}}}{r}$ is here $ut^{\frac{d+z-4}{z}}$, which again has to be compared to unity. Near the critical point, this correction is therefore negligible. However, a careful calculation of the terms arising when taking derivatives w.r.t. temperature shows that a nonanalyticity arises due to the temperature dependence of $r(t, \delta, u)$ in the classical regime. This leads to a divergence of the specific heat which is calculated in Appendix D.

In conclusion we find the variable u to give rise to corrections to coefficients in the limiting expansions of γ_{leading} (cf. (B2) and (B3), respectively) only – apart from a narrow region about the transition line $t_c(\delta)$, where the Gaussian correction to the discontinuity of the specific heat becomes important (see discussion in the last paragraph of Section II). The coefficients in the expansion of γ_{leading} are of order unity, whereas the corrections in $ur^{\frac{d+z-4}{2}}$ and $ut^{\frac{d+z-4}{z}}$ are small near the critical point.

APPENDIX D: SPECIFIC HEAT JUMP AND DIVERGENCE NEAR THE TRUE TRANSITION

The characteristic behavior of the model (2.2) near the transition is calculated in this section. As it has been shown in [3], (2.2) scales to a classical theory given by (2.22). For this theory, the discontinuity of the free energy reads [13]

$$\Delta F(b) = t(b) \frac{\delta(b)^2}{4v(b)} = \frac{\delta(b)^2}{4u(b)} \quad (\text{D1})$$

Using the relation $F(1) = b^{-(d+z)} F(b)$ in order to match the scaled model to the original one, we find the expression for the discontinuity of the free energy of model (2.2). We insert (2.6) for $u(b)$ and substitute $\delta(b) = b^2 r(t, \delta, u)$ with $r(t, \delta, u)$ given in Eq. (A10). The latter is

justified according to (2.11), noting the fact that the quantity $\Delta(t, \delta, u)$ becomes negligible in the classical regime. Differentiating twice w.r.t. temperature and taking into account that $r(t, \delta, u)$ vanishes at the transition, one ends up with the quantitative expression for the specific heat jump:

$$\Delta C = \mathcal{D}_{d,z}^2 N(N+2)^2 u t_c^{1+\frac{2}{z}(d-2)} \quad (\text{D2})$$

with the numerical constant $\mathcal{D}_{d,z}$ given by Eq. (A7). Please note the anomalous exponent of t_c in this expression. In theories where the distance to the critical point is measured by $(t - t_c)$ – classical Landau theories – we find that ΔC is proportional to t_c . For our case of a quantum critical point, the specific heat jump turns out to be smaller because additional (in general noninteger) powers of t_c occur.

Although the effective dimensionality of the theory (2.2) is $d + z$ (which is greater than 4 in the cases discussed in this communication), a divergence of the specific heat occurs very near the true transition (for vanishing $r(t, \delta, u)$). This is due to the fact that the quantum fluctuations become irrelevant and a classical description of the transition becomes more and more appropriate when the critical temperature is approached. To calculate the divergence, which has its origin in the temperature dependence of $r(t, \delta, u)$, we start from (2.18) and express the temperature derivatives (which we have to take in order to obtain γ) in terms of derivatives w.r.t. $r(t, \delta, u)$. Apart from contributions which are negligible corrections like those calculated in Appendix C, and analytic terms, we get

$$\gamma^{transition} = (\partial_t r)^2 \frac{2K_d N}{(2\pi)^{d+1}} r^{\frac{d+z-4}{2}} \int_0^\infty dy y^{\frac{4-d-z}{2}-1} \int_0^\infty d\varepsilon \coth\left(\frac{\varepsilon}{2\frac{t}{r^{z/2}}y^{\frac{z}{2}}}\right) \frac{\varepsilon (y+1)}{[\varepsilon^2 + (y+1)^2]^2} \quad (\text{D3})$$

The divergence occurs for $r(t, \delta, u) \rightarrow 0$, we can therefore expand the hyperbolic cotangent in its argument. Only the leading term gives rise to a divergence in $\gamma^{transition}$, which is

$$\gamma^{transition} = (\partial_t r)^2 \frac{4K_d N}{(2\pi)^{d+1}} t r^{\frac{d-4}{2}} \int_0^\infty dy y^{\frac{4-d}{2}-1} (y+1) \int_0^\infty d\varepsilon \frac{1}{[\varepsilon^2 + (y+1)^2]^2} \quad (\text{D4})$$

Performing the (elementary) integrals and using (A10) to evaluate $\partial_t r$ yields finally:

$$C^{transition} = \frac{NK_d(d-2)}{(2\pi)^{d-1}16\sin\frac{d-2}{2}\pi} \left(\frac{d+z-2}{z}\right)^{\frac{d-4}{2}} \left(\mathcal{D}_{d,z}(N+2)u t_c^{\frac{d+z-2}{z}}\right)^{\frac{d}{2}} \left[\frac{t-t_c}{t_c}\right]^{\frac{d-4}{2}} \quad (\text{D5})$$

The numerical constant $\mathcal{D}_{d,z}$ is given by Eq. (A7).

REFERENCES

- [1] for reviews see
G. R. Stewart, Reviews of Modern Physics **56** (1984) 755
C. M. Varma, Comments Solid State Physics **11** (1985) 221
P. A. Lee, T. M. Rice, J. W. Serene, L. J. Sham and J. W. Wilkins,
Comments Condensed Matter Physics **12** (1986) 99,
P. Fulde, J. Keller and G. Zwicknagel in *Solid State Physics*,
ed. by H. Ehrenreich and D. Turnbull, Vol. 41 Academic Press, New York 1988
- [2] C. J. Pethick and G. M. Carneiro, Physical Review **A7** (1973) 304
- [3] A. J. Millis, Physical Review **B48** (1993) 7183
- [4] M. A. Continentino, Physical Review **B47** (1993) 11587
- [5] A. M. Tsvelik, M. Reizer, Physical Review **B48** (1993) 9887
- [6] D. L. Cox, Physical Review Letters **59** (1987) 1240, J. Magn. Magn. Mater. **76-77**
(1988) 53
C. Sire, C. M. Varma, and H. R. Krishnamurthy, Physical Review **B48** (1993) 13833
and references therein
- [7] L. B. Ioffe and A. I. Larkin, Physical Review **B39** (1989) 8988
G. Baskaran, Z. Zou and P. W. Anderson, Solid State Commun. **63** (1987) 973
- [8] C. S. Jee et.al., Physical Review **B43** (1991) 2656
- [9] B. L. Altshuler, L. B. Ioffe, A. I. Larkin and A. J. Millis, unpublished
- [10] J. A. Hertz, Physical Review **B14** (1976) 1165
- [11] S. Sachdev, Zeitschrift für Physik **B94** (1994) 469
- [12] L. B. Ioffe and A. J. Millis, unpublished

- [13] S. K. Ma, Modern Theory of Critical Phenomena, Addison-Wesley 1976
- [14] D. R. Nelson, Physical Review **B11** (1975) 3504
- [15] L. D. Landau and E. M. Lifshitz, Statistical Physics Part I, Third edition
Pergamon Press, Oxford 1980
- [16] H. G. Schlager et.al., Journal of Low Temperature Physics **90** (1993) 181
- [17] A. Germann and H. v. Löhneysen, Europhys.Lett. **9** (1989) 367
- [18] A. deVisser et.al., JMMM **63&64** (1987) 365
- [19] A. deVisser, et.al., Physics Letters **113A** (1986) 489
- [20] G. E. Brodale et.al., Physical Review Letters **57** (1986) 234
- [21] F. Steglich et.al. Physical Review Letters **43** (1979) 1892

FIGURES

FIG. 1. Phase diagram for a quantum-critical point which is controlled by a zero temperature parameter δ . The shaded region is the (magnetically) ordered phase. The solid line denotes the transition curve $T_c(\delta)$, the dotted line corresponds to the quantum-classical crossover which occurs when T becomes of the order of $r^{z/2}$.

FIG. 2. Comparison of experimental data measured from CeCu_2Si_2 to the theoretical prediction for $d = 3, z = 2$. The temperature scale $T^* = 5\text{K}$ has been extracted from the data for $x = 0$, assuming the pure compound being at the critical point. The scale $x^* = 0.3$ is determined by the knowledge of T^* and the datapoint at $x = 0.05, T = 1.1\text{K}$. Solid curves represent the theoretical prediction, symbols are data points.